

UNIVERSITY OF NORTH CAROLINA
Department of Statistics
Chapel Hill, N. C.

Mathematical Sciences Directorate
Air Force Office of Scientific Research
Washington 25, D. C.

AFOSR Report No.

NOTE ON TRIANGULAR PARTIALLY BALANCED INCOMPLETE BLOCK DESIGNS

by

L. C. A. Corsten
Institute for Research on Varieties of Field Crops
Wageningen, The Netherlands

October, 1959

Contract No. AF 49(638)-213

The proper spaces of the matrix NN' , where N is the incidence matrix of triangular partially balanced incomplete block designs are exhibited explicitly in this paper. It also provides a convenient form of the Gramian of a basis of the join of two of these spaces.

Qualified requestors may obtain copies of this report from the ASTIA Document Service Center, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services, or have their "need-to-know" certified by the cognizant military agency of their project or contract.

Institute of Statistics
Mimeograph Series No. 242
October, 1959

NOTE ON TRIANGULAR PARTIALLY BALANCED INCOMPLETE BLOCK DESIGNS¹

by L. C. A. Corsten

Institute for Research on Varieties of Field Crops
Wageningen, The Netherlands

1. Summary.

The proper spaces of the matrix NN' where N is the incidence matrix of triangular partially balanced incomplete block designs are exhibited explicitly in this paper; it provides a convenient form of the Gramian of a basis of the join of two of these spaces.

2. Introduction.

The matrix N , the incidence matrix, for incomplete block designs is a matrix with v rows (v is the number of treatments) and b columns (b is the number of blocks) for which the typical element n_{ij} is unity if the i -th treatment occurs in the j -th block, and is zero otherwise. The non-negative symmetric matrix $Q=NN'$ of order v has elements q_{ij} , where q_{ii} is equal to the number of replicates of the i -th treatment, and q_{ij} ($i \neq j$) is equal to the number of blocks in which the i -th and the j -th treatment occur together.

In the case of partially balanced incomplete block designs with two associate classes, of which the class of triangular designs as developed by R. C. Bose and T. Shimamoto [1], is a subclass, the numbers

¹ This research was performed while the author was at the Department of Statistics of the University of North Carolina and was sponsored jointly by the United States Air Force through the Air Force Office of Scientific Research of the Air Research and Development Command under Contract No. AF 49(638)-213 and by a grant of the Netherlands Organization for Pure Research (Z.W.O.). Reproduction in whole or in part for any purpose of the United States Government is permitted.

q_{ii} are all equal to r (the number of replications), while the q_{ij} ($i \neq j$) are either λ_1 or λ_2 dependent on whether the pair of treatments i and j are first or second associates, respectively.

The knowledge of proper values and spaces of this Q is of interest in finding conditions of existence of designs with given sets of parameters; in addition it can contribute to a better understanding of the analysis of actually constructed designs and lead to the attachment of a physical meaning to their association schemes. We will pay attention to the last-mentioned points in a later paper.

The knowledge of the proper values of Q for several cases, including the triangular designs, as given by Connor and Clatworthy [2] has already been utilized for the derivation of necessary conditions for the parameters of such designs. The knowledge of the proper spaces of Q for triangular designs as will be shown in this note provided Ogawa [3] other conditions for existence of such designs.

3. The proper spaces of Q .

In order to consider the proper values and spaces of $Q=MN'$ we conceive Q as the matrix of the linear transformation Q of a vector space A consisting of vectors $\underline{x} = (x_1, x_2, \dots, x_v)$, into itself where the coordinate x_i corresponds to the i -th treatment.

As in partially balanced incomplete block designs with two associate classes the number of first and second associates of a fixed treatment is independent of the chosen fixed treatment, the i -th coordinate y_i in $\underline{y} = Q\underline{x}$ is equal to $rx_i + \lambda_1 S_1 + \lambda_2 S_2$, where S_j ($j=1,2$) represents the sum of the coordinates in \underline{x} corresponding to the j -th associates of treatment i .

We see immediately that if $\underline{z} = (1, 1, \dots, 1)$ then y_i is equal to $r + \lambda_1 n_1 + \lambda_2 n_2$ where n_j ($j=1, 2$) is the number of j -th associates of treatment i . As the n_j are independent of i in all partially balanced incomplete block designs it follows that $\underline{z} = (1, 1, \dots, 1)$ is a proper vector of Q with proper value $r + \lambda_1 n_1 + \lambda_2 n_2 = rk$ where k is the block size.

For further investigation of the proper spaces of Q we shall consider the subspace A^* of A orthogonal to \underline{z} . For every vector \underline{x} in A^* we have $x_i + S_1 + S_2 = 0$. Hence the coordinate y_i in $Q\underline{x}$ if \underline{x} is restricted to A^* is equal to

$$(1) \quad (r - \lambda_2)x_i + (\lambda_1 - \lambda_2)S_1.$$

For triangular designs the first and second associates of a treatment can be read from an association scheme which is constructed as follows. Consider a square array n by n in which the diagonal positions are empty (denoted by $*$) and where the remaining $n(n-1)$ positions each contain one of the $\frac{1}{2}n(n-1)$ treatment indices such that each index occurs twice and symmetrically with respect to the diagonal. For $n = 5$, e.g. this might be

$$\begin{pmatrix} * & 1 & 2 & 3 & 4 \\ 1 & * & 5 & 6 & 7 \\ 2 & 5 & * & 8 & 9 \\ 3 & 6 & 8 & * & 10 \\ 4 & 7 & 9 & 10 & * \end{pmatrix}$$

The first associates of any treatment are all those treatments which occur in the same row or the same column as this treatment, while the second associates are those which do not occur in the same row or the same column as this treatment. We note that $n_1 = 2(n-2)$ in this case.

For convenience we write the coordinates of any vector \underline{x} in A in the same arrangement as the corresponding indices in the upper diagonal part of the association scheme. We now construct a set of n vectors $\underline{c}_1, \underline{c}_2, \dots, \underline{c}_n$ in A in the following way. Write unity in all the positions of which the corresponding indices occur in the p -th row of the association scheme; write zero everywhere else; the resulting vector is called \underline{c}_p .

In our example with $n = 5$ we obtain

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & & \\ & & & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ & 1 & 1 & 1 \\ & & 0 & 0 \\ & & & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ & 1 & 0 & 0 \\ & & 1 & 1 \\ & & & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 & 0 \\ & 0 & 1 & 0 \\ & & 1 & 0 \\ & & & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 1 \\ & 0 & 0 & 1 \\ & & 0 & 1 \\ & & & 1 \end{pmatrix},$$

representing $\underline{c}_1, \dots, \underline{c}_5$, respectively.

Let the subspace of A of rank n spanned by these n linearly independent vectors be called A_1 . We note that A_1 contains the subspace spanned by \underline{z} . Now consider the inner product of any vector in A_1 :

$$\gamma_1 \underline{c}_1 + \dots + \gamma_n \underline{c}_n, \text{ and } \underline{z}; \text{ this is equal to } \gamma_1 (\underline{c}_1, \underline{z}) + \dots + \gamma_n (\underline{c}_n, \underline{z}) = (n-1) (\gamma_1 + \dots + \gamma_n). \text{ Hence for any vector in } A_1^*, \text{ the } (n-1)\text{-dimensional subspace of } A_1 \text{ orthogonal to } \underline{z}, \text{ we have: } \gamma_1 + \dots + \gamma_n = 0.$$

We further note that the coordinate of a vector in A_1 corresponding to the p -th row and the q -th column of the association scheme is equal to $\gamma_p + \gamma_q$.

Let \underline{x} be a vector in A_1^* . Then the sum of the coordinates of \underline{x} corresponding to all the treatments in the row and the column of the association scheme in which treatment i occurs (treatment i is counted twice in this sum) is equal to $2x_i + S_1$ on the one hand; on the other hand, according to the last two paragraphs, it is equal to:

$$\left\{ (n-1)\gamma_p + (\gamma_1 + \dots + \gamma_n) - \gamma_p \right\} + \left\{ (n-1)\gamma_q + (\gamma_1 + \dots + \gamma_n) - \gamma_q \right\} \\ = (n-2)(\gamma_p + \gamma_q) = (n-2)x_i,$$

if the treatment i occur in the association scheme in the p -th row and q -th column. Hence $S_1 = (n-4)x_i$ for all vectors in A_1^* .

Now it follows from (1) that the coordinate y_i of Qx where x is restricted to A_1^* is equal $\left\{ r + (n-4)\lambda_1 - (n-3)\lambda_2 \right\} x_i$. Therefore A_1^* is a proper space of Q with proper value $\left\{ r + (n-4)\lambda_1 - (n-3)\lambda_2 \right\}$.

Finally we consider the complement of A orthogonal to A_1 (which of course is the same as the complement of A^* orthogonal to A_1^*) and call this A_2 . The rank of A_2 is $\frac{1}{2}n(n-1) - n = \frac{1}{2}n(n-3)$.

Since every vector in A_2 is orthogonal to the given n basis vectors of A_1 , the sum of its coordinates corresponding to a row (or a column) of the association scheme must be zero. Taking the sum of its coordinates corresponding to the row and the column of the association scheme in which treatment i occurs (treatment i is counted twice again) we find: $2x_i + S_1 = 0$.

Again from (1) it then follows that the coordinate y_i of Qx where x is now restricted to A_2 is equal to: $(r - 2\lambda_1 + \lambda_2)x_i$. Therefore A_2 is also a proper space of Q and the corresponding proper value is $(r - 2\lambda_1 + \lambda_2)$.

4. The Gramian of A_1 .

For purposes of conditions for constructibility one is interested (see Ogawa [3]) in the Gramian, the symmetric matrix of inner products of a set of basis vectors of proper spaces of Q or of joins of such spaces. In this case it is quite easy to find the Gramian of the given

basis of A_1 , the join of the proper space A_1^* and the proper space spanned by \underline{z} . We simply need the inner products of the vectors $\underline{c}_1, \dots, \underline{c}_n$.

The diagonal elements of this Gramian are all equal to the number of unities in these basis vectors, i.e. $(n-1)$, while the off-diagonal elements are equal to the number of treatments which two different rows of the association scheme have in common, namely 1.

References

- [1] R. C. Bose and T. Shimamoto, "Classification and analysis of partially balanced incomplete block designs with two associate classes." J. Amer. Stat. Assn., Vol. 47 (1952), pp. 151-184.
- [2] W. R. Connor and W. H. Clatworthy, "Some theorems for partially balanced designs." Ann. Math. Stat., Vol. 25 (1954), pp. 100-112.
- [3] J. Ogawa, "A necessary condition for existence of regular and symmetrical experimental designs of triangular type, with partially balanced incomplete blocks." To be published.