

A RENEWAL THEOREM FOR SUMS OF I.I.D. VECTORS

by

Robert W. Keener

University of North Carolina at Chapel Hill

Abstract

The theorem we prove gives an approximation for the expected number of visits a drifting $(n+1)$ -dimensional random walk makes to a given set.

Running Head: Renewal Theory

AMS 1970 Subject Classifications: Primary 60K05; Secondary 60J15.

Key Words and Phrases: renewal theory, random walks.

This research was supported by National Science Foundation Grant MCS78-01240 and Air Force Office of Scientific Research Grant AFOSR-80-0080.

Although a great deal of research has been devoted to renewal limit theorems in one dimension, the literature concerning renewal theory in several dimensions is rather sparse. Doney [4] and Stam [11] have studied the asymptotic renewal measure of a translated set and Mode [8] has derived a related renewal density theorem.

In our theorem we obtain a finite asymptotic limit for the renewal measure of a set as it is translated in the direction of the mean and scaled in the other directions. The theorems of Doney, Stam and Mode are related to our result much as the local limit theorem is related to the central limit theorem. Other limit theorems in the same general area have been given by Smith [10], Bickel and Yahav [1], Farrell [5,6], and Ney and Spitzer [9].

Let X be a random variable and Y an n -dimensional random vector, with joint law P . Define the renewal measure

$$U = \sum_{i=0}^{\infty} P^{(i)}$$

where $P^{(i)}$ is the i -fold convolution of P with itself. For a set $S \in \mathbb{R}^{n+1}$ let

$$S_{\tau} = \{(x,y) : (x,y/\tau) \in S\}.$$

The following result is a multi-dimensional generalization of Blackwell's [2,3] renewal theorem.

Theorem. If X is non-lattice, has a finite second moment and a finite positive mean μ , and if Y has zero mean and finite covariance $\Sigma = EYY'$, then for any Borel set S whose projection onto the x -axis is bounded we have

$$\lim_{\tau \rightarrow \infty} U((\tau^2, 0) + S_\tau) = \int_S \frac{dx}{\mu} N(0, \Sigma/\mu)(dy)$$

where $N(\cdot, \cdot)$ is the multivariate normal measure.

Proof. For constants $b = 0, 1$ and $r < 1$, define the measures V_τ by

$$V_\tau(S) = \sum_{i=0}^{\infty} r^i [P^{(i)}((b\tau^2, 0) + S_\tau) + P^{(i)}((-b\tau^2, 0) - S_\tau)] .$$

If ϕ is the characteristic function

$$\phi(p, q) = E \exp(ipX + iq'Y) ,$$

then the Fourier transform of V_τ is

$$\int e^{ipx+iq'y} V_\tau(dx, dy) = 2e^{-ipb\tau^2} \operatorname{Re} \left[\frac{1}{1-r\phi(p, q/\tau)} \right] .$$

If g is an L^1 function such that

$$\hat{g}(z) = \int g(x) e^{-izx} dx$$

is L^1 , then Parseval's relation gives

$$(1) \quad \int g(x) e^{iq'y} V_\tau(dx, dy) = \frac{1}{\pi} \int \hat{g}(z) e^{-ib\tau^2 z} \operatorname{Re} \left[\frac{1}{1-r\phi(z, q/\tau)} \right] dz .$$

If we choose

$$g(x) = \frac{3}{2h} \left(1 - \frac{|x|}{h}\right)^2 \quad \text{for } |x| < h$$

$$= 0 \quad \text{otherwise}$$

and $b = 0$, we have

$$\frac{3}{2h} \int_{-h}^h \left(1 - \frac{|x|}{h}\right)^2 e^{iq'y} V_{\tau}(dx, dy) = \frac{6}{\pi h^2} \int \left(1 - \frac{\sin(zh)}{zh}\right) \operatorname{Re} \left[\frac{1}{1 - r\phi(z, q/\tau)} \right] \frac{dz}{z^2},$$

which implies

$$\frac{2}{h} \int_{|z| > \frac{2}{h}} \operatorname{Re} \left[\frac{1}{1 - r\phi(z, q/\tau)} \right] \frac{dz}{1+z^2} < \pi \lim_{r \rightarrow 1} V_{\tau}((-h, h) \times \mathbb{R}^n).$$

Since X has a positive mean, this last expression is uniformly bounded for positive τ ; so as $h \downarrow 0$,

$$(2) \quad \int_{|z| > \frac{2}{h}} \operatorname{Re} \left[\frac{1}{1 - r\phi(z, q/\tau)} \right] \frac{dz}{1+z^2} = O(h)$$

uniformly in τ , q and r . For the rest of our argument we will take $b = 1$,

$$g(x) = e^{ipx} / (\pi(1+x^2)),$$

$$\hat{g}(z) = e^{-|z-p|},$$

and let $r < 1$ be a function of τ , whose difference from 1 is $o(1/\tau^2)$ as $\tau \rightarrow \infty$. From Taylor's Theorem, if $z \rightarrow 0$ and $\tau \rightarrow \infty$ in a manner such that $z = o(1/\tau^2)$,

$$(3) \quad \operatorname{Re} \left[\frac{1}{1-r\phi(z, q/\tau)} \right] = \frac{2\tau^2 q' \Sigma q}{4\mu^2 z^2 \tau^4 + (q' \Sigma q)^2} + o(\tau^2),$$

and if $z \rightarrow 0$ and $\tau \rightarrow \infty$ in a manner such that $1/\tau^2 = o(z)$,

$$(4) \quad \operatorname{Re} \left[\frac{1}{1-r\phi(z, q/\tau)} \right] = O \left(1 + \frac{1}{\tau^2 z^2} \right).$$

We now fix p, q and $\epsilon > 0$. Feller and Orey [7] have shown that

$$\frac{1}{z^2 + 1} \operatorname{Re} \left[\frac{1}{1-\phi(z, 0)} \right]$$

is L^1 . Using this fact and equations (2) and (4) it is possible to choose

$\delta = 1/M$ so that for $\tau > M$,

$$\left[\int_{|z| > M} + \int_{|z| < \delta} \right] \frac{\hat{g}(z)}{\pi} \operatorname{Re} \left[\frac{1}{1-\phi(z, 0)} \right] dz \leq \frac{\epsilon}{6},$$

$$\left[\int_{|z| > M} + \int_{\frac{M}{\tau^2}}^{\delta} + \int_{-\delta}^{-\frac{M}{\tau^2}} \right] \frac{\hat{g}(z)}{\pi} \operatorname{Re} \left[\frac{1}{1-r\phi(z, q/\tau)} \right] dz \leq \frac{\epsilon}{6},$$

and

$$\int_{|z| > M} \frac{\hat{g}(0)}{\pi} \frac{2q' \Sigma q}{4\mu^2 t^2 + (q' \Sigma q)^2} dt \leq \frac{\epsilon}{6}.$$

Using the Riemann-Lesbegue Lemma, equation (3) and the fact that X is non-lattice, we see that for τ sufficiently large we will have

$$\left| \int \frac{\hat{g}(z)}{\pi} e^{-iz\tau^2} \operatorname{Re} \left[\frac{1}{1-\phi(z,0)} \right] dz \right| \leq \frac{\varepsilon}{6} ,$$

$$\left| \frac{1}{\pi} \int_{-\frac{M}{\tau^2}}^{\frac{M}{\tau^2}} \left\{ \hat{g}(z) e^{-iz\tau^2} \operatorname{Re} \left[\frac{1}{1-r\phi(z,q/\tau)} \right] - \hat{g}(0) e^{-iz\tau^2} \frac{2\tau^2 q' \Sigma q}{4\mu^2 z^2 \tau^4 + (q' \Sigma q)^2} \right\} dz \right| \leq \frac{\varepsilon}{6} ,$$

and

$$\left| \left\{ \int_{\delta}^M + \int_{-M}^{-\delta} \right\} \frac{\hat{g}(z)}{\pi} e^{-iz\tau^2} \operatorname{Re} \left[\frac{1}{1-r\phi(z,q/\tau)} - \frac{1}{1-\phi(z,0)} \right] dz \right| \leq \frac{\varepsilon}{6} .$$

Combining the last 6 inequalities we see that

$$\left| \int \frac{\hat{g}(z)}{\pi} e^{-iz\tau^2} \operatorname{Re} \left[\frac{1}{1-r\phi(z,q/\tau)} \right] dz - \frac{\hat{g}(0)}{\pi} \int \frac{2q' \Sigma q}{4\mu^2 t^2 + (q' \Sigma q)^2} e^{-it} dt \right| < \varepsilon .$$

As ε was arbitrary, this proves that

$$\begin{aligned} \lim_{\tau \rightarrow \infty} \int \frac{e^{ipx+iq'y}}{\pi(1+x^2)} V_{\tau}(dx, dy) &= \frac{1}{\mu} \exp \left[-\frac{q' \Sigma q}{2\mu} - |p| \right] \\ &= \int \frac{e^{ipx+iq'y}}{\pi(1+x^2)} \frac{dx}{\mu} N(0, \Sigma/\mu)(dy) . \end{aligned}$$

As the rate at which $r(\tau) \rightarrow 1$ was arbitrary, this equation must also hold with $r = 1$, and by the Lévy Continuity Theorem, the laws with density $1/(\pi(1+x^2))$ with respect to V_{τ} converge weakly. Hence, if S is a Borel set whose projection onto the x -axis is bounded we have

$$\lim_{\tau \rightarrow \infty} \{U((\tau^2, 0) + S_{\tau}) + U((-\tau^2, 0) - S_{\tau})\} = \int_S \frac{dx}{\mu} N(0, \Sigma/\mu)(dy) .$$

This completes the proof as

$$\lim_{\tau \rightarrow \infty} U((-\tau^2, 0) - S_\tau) = 0 .$$

References

- [1] Bickel, P.J. and Yahav, J.A. (1965). Renewal theory in the plane. *Ann. Math. Statist.* 36, 946-955.
- [2] Blackwell, D.H. (1948). A renewal theorem. *Duke Math J.* 15, 145-150.
- [3] Blackwell, D.H. (1953). Extension of a renewal theorem. *Pacific J. Math.* 3, 315-320.
- [4] Doney, R.A. (1966). An analogue of the renewal theorem in higher dimensions. *Proc. London Math. Soc., Ser. 3* 16, 669-684.
- [5] Farrell, R.H. (1964). Limit theorems for stopped random walks. *Ann. Math. Statist.* 35, 1332-1343.
- [6] Farrell, R.H. (1966). Limit theorems for stopped random walks III. *Ann. Math. Statist.* 37, 1510-1527.
- [7] Feller, W. and Orey, S. (1961). A renewal theorem. *J. Math. Mech.* 10, 619-624.
- [8] Mode, C.J. (1967). A renewal density theorem in the multi-dimensional case. *J. Appl. Prob.* 4, 62-76.
- [9] Ney, P. and Spitzer, F. (1966). The Martin boundary for random walks. *Trans. Am. Math. Soc.* 121, 116-132.
- [10] Smith, W.L. (1955). Regenerative stochastic processes. *Proc. Royal Soc. A* 232, 6-31.
- [11] Stam, A.J. (1969). Renewal theory in r dimensions (I). *Comp. Math.* 21, 383-399.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Renewal Theorem for Sums of I.I.D. Vectors		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
		6. PERFORMING ORG. REPORT NUMBER Mimeo Series No. 1286
7. AUTHOR(s) Robert W. Keener		8. CONTRACT OR GRANT NUMBER(s) NSF Grant MCS78-01240 AFOSR Grant AFOSR-80-0080
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE May 1980
		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release -- Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) renewal theory, random walks		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The theorem we prove gives an approximation for the expected number of visits a drifting (n+1)-dimensional random walk makes to a given set.		

